

Юрий Кузьменко

Переводы

Country	Украина	Phone	+380963460045
City	Винница	Email	did96117@gmail.com

Профессиональное резюме

FLIGHTS WITH THE AIRLINE FOR 15 YEARS ALLOWED ME TO IMPROVE THE KNOWLEDGE OF FOREIGN LANGUAGES, ESPECIALLY ENGLISH. THIS IS MY FAVORITE HOBBY FROM CHILDHOOD!

Трудовая биография

бортпроводник международных авиалиний	авиакомпания УзбекистанЭйрВэйз	Ташкент
2023-06-14 – 2023-06-14		

Профессиональные навыки

Мои переводы - Ampicillin is an antibiotic used to treat various infectious diseases of the urinary, respiratory tract, liver and gastrointestinal tract. It is ineffective against penicillinase, so it should not be given in cases where the disease is caused by microbes that produced by this enzyme. It is taken orally or injection is appointed, the possible side effects: nausea, vomiting, diarrhea, may develop various allergic reactions. Trade names: amfipen (Amfipen), penbritin (Penbritin).

The drug is not destroyed in the acidic environment of the stomach, it is well absorbed when taken orally. It is active against Gram-positive microorganisms, which operates benzyl penicillin. In addition, it operates on a number of gram-negative microorganisms (Salmonella, Shigella, Proteus, Escherichia coli, Klebsiella pneumonia (pneumobaccillus), Pfeiffer's bacillus (bacillus influenza)) and it is therefore seen as a broad spectrum antibiotic and is used in diseases caused by mixed infection.

Indications:

Infectious-inflammatory diseases caused by microorganisms susceptible to Ampicillin, including infections of the ear, nose and throat, Odontogenous infections, respiratory tract infections (pneumonia, bronchitis, pneumonia, lung abscess), acute and chronic urinary infections and biliary tract (pyelitis, pyelonephritis, cholecystitis), gastrointestinal infections (including salmonella), gynecological infections (including gonorrhoea), meningitis, endocarditis, septicemia, sepsis, rheumatic fever, erysipelas, scarlet fever, infections of the skin and soft tissue.

Contraindications:

Hypersensitivity (including to penicillin and other beta-lactam antibiotics), infectious mononucleosis, lymphocytic leukemia.

Pregnancy and breastfeeding:

During pregnancy, it is possible in case the expected effect of therapy exceeds a potential risk to the fetus. At the time of treatment by the drug, breastfeeding shall be going on, but the duration of treatment is minimized (it passes into breast milk).

Undesirable effects:

For the cardiovascular system and blood (hematopoiesis, hemostasis): anemia, leucopenia, thrombocytopenia, and agranulocytosis.

For the digestive tract: nausea, vomiting, diarrhea, dysbiosis, pseudo membranous colitis, glossitis, stomatitis, increased activity of hepatic transaminases.

Allergic reactions: skin rash, itching, urticaria, rhinitis, conjunctivitis, fever, arthralgia, eosinophilia, angioedema; exfoliate dermatitis, erythematic multiform, anaphylactic shock, etc.

Other: oral candidiasis, vaginal candidiasis.

Переводы

различная тематика.

совершенствование знаний с помощью общения с носителями языка из 26

26 стран, посещенных в полетах, где большой процент граждан владеет иностранными языками, в большинстве своем английским. Если бы люди всей земли лучше бы понимали друг друга, культуру, обычаи и особенности, было меньше конфликтов и больше практической пользы. В том числе это и есть одна из задач переводчиков.
